

## References (30 publications over 10 years, as background to vigour tests)

- Basak, O., Demir, I., Mavi, K. and Matthews, S. (2006). Controlled deterioration test for predicting seedling emergence and longevity of pepper (*Capsicum annuum* L.) seed lots. *Seed Science and Technology*, **34**, 723-734.
- Demir, I., Cebeci, C. and Guloksuz, T. (2012). Electrical conductivity measurements to predict germination of commercially available radish seed lots. *Seed Science and Technology*, **40**, 229-237.
- Demir, I., T. Celikkol, G. Sari Kumis and C. Eksi. 2011. Vigour tests to estimate seedling emergence potential and longevity in *Viola* seed lots. *HortScience*, **46**, 402-405.
- Demir, I., S. Ermis, K. Mavi and S. Matthews. 2008. Mean germination time of pepper seed lots (*Capsicum annuum* L.) predicts size and uniformity of seedlings in germination tests and transplant modules. *Seed Science and Technology*, **36**, 21-30.
- Demir, I., Mavi, K., Kenanoglu, B.B. and Matthews, S. (2008) Prediction of germination and vigour in naturally aged commercially available seed lots of cabbage (*Brassica oleracea* var *capitata*) using the bulk conductivity method. *Seed Science and Technology*, **36**, 509-523.
- Ermis, S., Karshloglu, M., Ozden, E. and Demir, I. (2015), Use of a single radicle emergence count as a vigour test in prediction of seedling emergence potential of leek seed lots *Seed Science and Technology*, **43**, 308-312.
- Ermis, S., Ozden, E. and Demir, I. (2015), Early radicle emergence count after controlled deterioration (CD) predicts emergence potential better than final standard germination after CD in tomato varieties *Seed Science and Technology*, **43**, 278-283.
- Ilbi H and Eser B . (2006). The potential of vigour tests to identify differences in the extent of ageing in onion seeds. *Seed Science and Technology*, **34**, 713-718.
- Ilbi, H., Kavak, S. and Eser, B. (2009). *Seed Science and Technology*, **37**, 516-519. Cool germination test can be an alternative vigour test for maize
- Kavak, S., Ilbi, H. and Eser, B. (2008). Controlled deterioration test determines vigour and predicts field emergence in pepper seed lots. *Seed Science and Technology*, **36**, 456-461.
- Khajeh Hosseini, M., A. Lomholt and S. Matthews. (2009). Mean germination time in the laboratory estimates the relative vigour and field performance of commercial seed lots of maize (*Zea mays* L). *Seed Science and Technology*, **37**, 446-456.
- Khajeh Hosseini, M., M. Nasehzadeh and S. Matthews. 2010. Rate of physiological germination relates to the percentage of normal seedlings in standard germination tests of naturally aged seed lots of oil seed rape. *Seed Science and Technology*, **38**, 602-611.
- Lv, Y.Y., Wang, Y.R. and Powell, A.A. (2016). Frequent individual counts of radicle emergence and mean just germination time predict seed vigour of *Avena sativa* and *Elymus nutans*. *Seed Science and Technology*, **44**, 189-198.
- Matthews, S., E. Beltrami, R. El-Khadem, M. Khajeh Hosseini, M. Nasehzadeh and G. Urso. (2011). Evidence that time for repair during early germination leads to vigour differences in maize. *Seed Science and Technology*, **39**, 501-509.

- Matthews, S., Demir, I., Celikkol, T., Kenanoglu, B.B. and Mavi, K. (2009), Vigour tests for cabbage seeds using electrical conductivity and controlled deterioration to estimate relative emergence in transplant modules. *Seed Science and Technology*, **37**, 736-746
- Matthews, S., R. El-Khadem, E. Casarini, M. Khajeh Hosseini, M. Nasehzadeh and M.-H. Wagner. 2010. Rate of physiological germination compared with the cold test and accelerated ageing as a repeatable vigour test for maize (*Zea mays*). *Seed Science and Technology*, **38**, 379-389.
- Matthews, S. and Khajeh-Hosseini, M. (2006). Mean germination time as an indicator of emergence performance in soil of seed lots of maize (*Zea mays*) *Seed Science and Technology*, **34**, 361-369.
- Matthews, S. and Khajeh Hosseini, M. 2007. Length of the lag period of germination and metabolic repair explain vigour differences in seed lots of maize (*Zea mays*). *Seed Science and Technology*, **35**, 200-212.
- Matthews, S., Noli, E., Demir, I., Khajeh-Hosseini, M. and Wagner, M-H. (2012). Evaluation of seed quality: from physiology to international standardisation. *Seed Science Research*, **22**, S69-S73.
- Matthews, S. and Powell, A.A. (2011). Towards automated single counts of radicle emergence to predict seed and seedling vigour. *Seed Testing International*, **142**, 44-48.
- Matthews, S., Wagner M.-H, Ratzenboeck A., Khajeh Hosseini M., Casarini E., El-Khadem R., Yakhlifi M. and Powell A.A.. (2011). Early counts of radicle emergence during germination as a repeatable and reproducible vigour test for maize. *Seed Testing International*, **141**, 39-45.
- Matthews S., Wagner M.-H., Kerr L., McLaren G. and Powell A.A. (2012). Automated determination of germination time courses by image capture and early counts of radicle emergence lead to a new vigour test for winter oilseed rape (*Brassica napus*) *Seed Science and Technology*, **40**, 413-424.
- Mavi, K., Demir, I. and Matthews, S. (2010). Mean germination time estimates the relative emergence of seed lots of three cucurbit crops under stress conditions. *Seed Science and Technology*, **38**, 14-25.
- Mavi, K., Mavi, F., Demir, I. and Matthews, S. (2014). Electrical conductivity of seed soak water predicts seedling emergence and seed storage potential in commercial seed lots of radish. *Seed Science and Technology*, **42**, 76-86.
- Mériaux B., Wagner M.- H., Ducournau S., Ladonne F., Fougereux J.- A. (2007). Using sodium chloride saturated solution to standardize accelerated ageing test for wheat seeds. *Seed Science and Technology* **35**, 722-732.
- Mirdad Z, Powell, A.A. and Matthews, S. (2006). Prediction of germination in artificially aged seeds of *Brassica* spp. using the bulk conductivity test. *Seed Science and Technology*, **34**, 273-286.
- Noli, E., Casarini, G., Urso, G. and Conti, S. (2008). Suitability of three vigour test procedures to predict field performance of early sown maize seed. *Seed Science and Technology*, **36**, 168-176.
- Noli E, Beltrami E, Casarini E, Urso G, Conti S (2010) Reliability of early and final counts in cold and cool germination tests for predicting maize seed vigour. *Italian Journal of Agronomy*, **4**, 383–391.

Powell, A.A. and Matthews, S. (2012). Seed aging/repair hypothesis leads to new testing methods. *Seed Technology*, **34**, 15-25

Wagner M.H. and Ducournau S. (2007). Conductivity testing for oilseed rape seeds. *Seed Testing International*, **133**, 40-41.

In addition, although not vigour, its basis has arisen out of vigour work

Mavi, K., Powell, A.A. and Matthews, S. (2016). Rate of radicle emergence and leakage of electrolytes provide quick predictions of percentage normal seedlings in standard germination tests of radish (*Raphanus sativus*). *Seed Science and Technology*, **44**, in press.

### **Validation reports (11 in 10 years: in chronological order to show progress in vigour test method development)**

All can be found at <http://www.seedtest.org/en/method-validation-reports-content---1--1256.html>

Powell, A.A. (2006). Proposal for the addition of *Phaseolus vulgaris* as a species to which the conductivity test for seed vigour can be applied.

Powell, A.A. (2008). Evaluation of the controlled deterioration test as a repeatable and reproducible vigour test for Brassica species

Powell, A.A. (2009). Proposal for the addition of *Glycine max* as a species to which the conductivity test for seed vigour can be applied.

Matthews S., Wagner M-H., Ratzenboeck, A., Khajeh-Hosseini, M., Casarini, E., El-Khadem, El Yakhli, M. and Powell, A.A. (2011). Evaluation of early counts of radicle emergence during germination as a repeatable and reproducible vigour test for maize

Khajeh-Hosseini, M., Gallo, H. and Ilbi, H. (2014). Proposal for the addition of *Cicer arietinum* (Kabuli type) as a species to which the conductivity test for seed vigour can be applied.

Powell, A.A., Matthews, S., Kerr, L., McLaren, G. and Wagner M-H. (2014) Early counts of radicle emergence as a vigour test for oil seed rape.

Gallo, C.; França-Neto, J.B; Arango, M.; Gonzalez, S.; Francomano, V.; Carracedo, C.; Costa, O.; Alves, R.; Craviotto, R. (2016). Proposal for the addition of tetrazolium method as a vigour test for *Glycine max* seeds. <http://www.seedtest.org/upload/cms/user/OGM16-06-Method-Validation-Reports-for-Rules-2017.pdf>.

Powell, A.A. and Mavi, K. (2016). Application of the electrical conductivity test to radish seed (*Raphanus sativus*). Method validation report, International Seed Testing Association. <http://www.seedtest.org/upload/cms/user/OGM16-06-Method-Validation-Reports-for-Rules-2017.pdf>

Powell, A.A. and Mavi, K. (2016). Application of the radicle emergence test to radish (*Raphanus sativus*) seed). Method validation report, International Seed Testing Association. <http://www.seedtest.org/upload/cms/user/OGM16-06-Method-Validation-Reports-for-Rules-2017.pdf>

Wagner M-H., Powell, A.A., Corre, F., McLaren, G., Noli, E., Matthiessen, C. and Ducournau, S. (2016) Alternative method for seed moisture content adjustment in vigour testing, as applied in the CD

test. <http://www.seedtest.org/upload/cms/user/OGM16-06-Method-Validation-Reports-for-Rules-2017.pdf>.

Wagner M-H., Powell, A.A., El-Yakhlifi, M. and Ducournau, S. (2016) Shortening the controlled deterioration (CD) test for Brassica by replacing the germination test with a conductivity measurement. <http://www.seedtest.org/upload/cms/user/OGM16-06-Method-Validation-Reports-for-Rules-2017.pdf>.